

Rev. Inst. Med. trop. S. Paulo
51(4):219-221, July-August, 2009
doi: 10.1590/S0036-46652009000400008

BRIEF COMMUNICATION

***Ascaris lumbricoides*: ALTERATION OF THE ERYTHROCYTE SUPERFICIAL CHARGE USING THE PARTITION METHOD IN AQUEOUS TWO-PHASE SYSTEM**

Patricia PONCE DE LEÓN, Natalia LEBENSOHN, Patricia FORESTO & Juana VALVERDE

SUMMARY

Sialic acid is responsible for the negative charge of the erythrocyte. The decrease of sialic acid has hemodynamical and hemorheological importance. The aim was to study the effect of *A. lumbricoides* on the erythrocyte superficial charge using the Partition Method in aqueous two-phase system in order to indirectly evaluate the alteration of sialic acid in the red cells.

We worked with five parasite extracts (AE) and larvae concentrate (LC). Erythrocyte superficial charge was studied by working with non-treated (Controls) and treated erythrocytes. The treatment consisted of incubating the erythrocytes with AE or LC for 30 minutes at 4 °C, 20 °C and 37 °C. The red cells were separated in a sensitive charge two-phase system (Dx/ PEG). The partition coefficient (P) of treated and untreated erythrocytes were calculated. The results showed a P decrease at the three temperatures for red cells treated with four of the AE. The remaining extract did change P values at any of the temperatures studied. The erythrocytes treated with LC showed a decrease in the P value at 37 °C and 4 °C but no change was observed at 25 °C. Statistical analysis concluded that P values were significantly lower in treated erythrocytes than in their corresponding untreated ones ($p < 0.05$). The Partition Method showed that this parasite alters the erythrocyte superficial charge which may indicate that it can catch sialic acid.

KEYWORDS: *Ascaris lumbricoides*; Erythrocyte charge; Partition method.

Sialic acid is responsible for the negative charge of the erythrocyte membrane and contributes to the structural properties of the glycoproteins. A decrease in the contents of sialic acid in the membrane promotes erythrocyte aggregation, rouleaux formation, decrease in the blood flow and increase in blood viscosity, which results in a greater interaction between erythrocytes and vascular endothelium⁸. Cellular mechanisms of many infectious processes are not known with accuracy and they may involve host and parasite surface glycoconjugates. *Trypanosoma cruzi* takes sialic acid from the host glycoconjugates and transfers it to molecules on its surface. Taking hold of sialic acid makes *T. cruzi* resistant to the attack of complement proteins and antibodies of cytolytic capacity⁵. In trypanosomiasis caused by *T. congolense*, *T. brucei* and *T. evansi* the adhesion phenomenon of the parasites to erythrocytes and endothelial cells is mediated by structures which contain sialic acid residues^{2,4,9}. The existence of invasion ways through acid sialic receptors in *Plasmodium falciparum* has been communicated⁷. The occurrence of sialic acid in different parasite infections with endocellular replication has been reported but there are no reports of its role in *Ascaris lumbricoides*. The aim was to study the effect of *A. lumbricoides* on the erythrocyte superficial charge using the Partition Method (PM) in aqueous two-phase system in order to indirectly evaluate the alteration of sialic acid in the red cells.

Five parasite extracts (AE) and a larvae concentrated (LC) (500-600 larvae/mL) were used. AE were obtained by surgical remotion of the cuticle from adult worms¹³ and LC was obtained from eggs eclosion⁶ and larvae collection by Baerman Morales Method¹⁶. Erythrocyte superficial charge was studied by working with non -treated human erythrocytes (controls) and treated (T) ones, both washed in phosphate buffer saline (PBS) and resuspended to a 75% hematocrit. The red cells were obtained from normal individuals O Group. The treatment consisted of incubating the erythrocytes with AE or LC for 30 minutes at 4 °C, 20 °C and 37 °C. Then, they were washed three times with PBS. The red cells were separated in an aqueous system containing 5% (w/w) Dx and 4 % (w/w) PEG in PBS, pH 7.4. The erythrocytes are retained of the upper phase according to the negative superficial charge^{1,15}. The partition coefficient (P) is defined as the number of cells in the upper phase with respect to the totality of added cells. P of untreated erythrocytes was compared to the P obtained from the same treated red cells. The P values were simultaneously calculated in treated and untreated erythrocytes. Fifteen measurements were made for the 5 AE at the three temperatures and six measurements were made for LC which was incubated in two erythrocyte suspensions at each temperature.

25 °C: PNT = 0.91 / PT [EA] ₁ = 0.15	4°C: PNT = 1 / PT [EA] ₁ = 0.31	37 °C: PNT = 1 / PT [EA] ₁ = 0.60
PNT = 0.91 / PT [EA] ₂ = 0.08	PNT = 0.92 / PT [EA] ₂ = 0.56	PNT = 1 / PT [EA] ₂ = 0.43
PNT = 1 / PT [EA] ₃ = 0.27	PNT = 0.92 / PT [EA] ₃ = 0.17	PNT = 1 / PT [EA] ₃ = 0.17
PNT = 0.91 / PT [EA] ₄ = 0.50	PNT = 0.92 / PT [EA] ₄ = 0.18	PNT = 1 / PT [EA] ₄ = 0.59
PNT = 1 / PT [EA] ₅ = 0.97	PNT = 0.96 / PT [EA] ₅ = 0.90	PNT = 0.90 / PT [EA] ₅ = 0.90
PNT = 0.95 / PT CL = 0.95	PNT = 1 / PT CL = 0.02	PNT = 0.98 / PT CL = 0.01
PNT = 0.98 / PT CL = 0.98	PNT = 1 / PT CL = 0.01	PNT = 0.98 / PT CL = 0.02

PNT: value of the partition coefficient of non- treated erythrocytes; PT: value of the partition coefficient of treated erythrocytes

The results showed a decrease in P value at the three temperatures for erythrocytes treated with four of the five AE. The remaining AE did not modify P values at any of the temperatures assayed. The erythrocytes treated with LC showed a decrease in P value at 37 °C and 4 °C. No change was observed at 25 °C. The observed P values were lower in red cells treated with LC than in erythrocytes treated with AE.

In order to compare PT values and their respective PNT, Wilcoxon Test¹⁷ was used in paired samples for each temperature. The results of statistical analysis concluded that P values were significantly lower in treated erythrocytes than in their corresponding untreated ones ($p < 0.05$). At 4 °C PNT median was 0.96 and the interquartile range was 0.08 while PT values were 0.18 and 0.54, respectively. At 25 °C, the P median for the controls was 0.95 and the interquartile range 0.09, while PT were 0.50 and 0.82, respectively. At 37 °C, PNT median was 1 and the interquartile range 0.02, while PT values were 0.43 and 0.58, respectively. P values showed a greater variation in treated red cells than in untreated ones at the three temperatures.

PM is a simple technique used to separate cells having slightly different surface properties and to characterize biologically important molecules^{1,15}. Authors of this experience have used PM to study changes in erythrocyte superficial charge^{8,10,11,12,14}. The results have shown that *A. lumbricoides* can alter the erythrocyte superficial charge which may indicate that it can catch sialic acid both in its adult and larval stages. The larvae stages circulate into the blood stream. The adult stage, even if it is located in intestine, might have contact with blood when it pierces the intestinal wall or penetrates into ducts and veins³. This finding could be significant to explain the evading mechanism of the host's immune response.

RESUMEN

Ascaris lumbricoides: alteración de la carga superficial eritrocitaria usando el Método de Partición en sistemas bifásicos acuosos

La disminución de ácido sialico, responsable de la carga negativa del eritrocito, tiene importancia hemodinámica y hemorreológica. El objetivo fue estudiar el efecto de *A. lumbricoides* sobre la carga superficial eritrocitaria aplicando el método de partición en sistemas bifásicos acuosos, a los fines de evaluar de manera indirecta la alteración de ácido sialico de los eritrocitos. Se trabajó con 5 extractos del parásito adulto (EA) y con un concentrado de larvas (500-600 larvas/mL) (CL). Se estudió la carga superficial eritrocitaria, trabajando con eritrocitos no tratados y tratados. El tratamiento consistió en la incubación de los eritrocitos con EA o CL durante 30 minutos a 4 °C, 25 °C y 37 °C. Los eritrocitos fueron sometidos a la separación en un sistema bifásico carga

sensible constituido por Dx / PEG. Se calculó el coeficiente de reparto (P), de los eritrocitos sin tratar y tratados. Los resultados mostraron disminución de P a las 3 temperaturas, en hematíes tratados con 4 de los EA. El EA restante no modificó los valores de P a ninguna de las temperaturas estudiadas. CL produjo la disminución de P a 37 °C y 4 °C, pero no se observó modificación a 25 °C. Los análisis estadísticos concluyeron que los valores de P son significativamente menores en los eritrocitos tratados que en los respectivos eritrocitos sin tratar ($p < 0.05$). El método de partición demostró que *A. lumbricoides* altera la carga superficial eritrocitaria lo que indicaría que el parásito, tanto en su estado adulto como en sus fases larvales, puede captar ácido sialico.

REFERENCES

- ALBERTSON, P.A. - **Partition of cell particles and macromolecules: distribution and fractionation of cells, mitochondria, chloroplasts, viruses, proteins, nucleic acids, and antigen-antibody complexes in aqueous polymer two-phase systems**. New York, Wiley-Interscience, 1971.
- ANOSA, V.O. & KANEKO, J.J. - Pathogenesis of *Trypanosoma brucei* infection in deer mice (*Peromyscus maniculatus*): light and electron microscopic studies on erythrocyte pathologic changes and phagocytosis. **Amer. J. vet. Res.**, 44: 645-651, 1983.
- BEAVER, P.C.H.; JUNG, R.C. & CUPP, E.W. - **Parasitología clínica**. Barcelona, Salvat, 1986.
- BOADA SUCRE, A.; ROSSI, S.; MARCELLO, S. & DE STEFANO, H. - Alteraciones en el patrón de enlazamiento de lectinas en el riñón de ratones infectados experimentalmente con una cepa venezolana de *Trypanosoma evansi*. **Rev. Cient. (Maracaibo)**, 14: 431-439, 2004.
- BUSCAGLIA, C. - Trans-sialidasa de *Trypanosoma cruzi*: un blanco potencial para el tratamiento de la enfermedad de Chagas. **Rev. Hosp. Matern. infant. Ramón Sardá**, 21: 24-27, 2002.
- FAIRBAIRN, D. - The *in vitro* hatching of *Ascaris lumbricoides* eggs. **Canad. J. Zool.**, 39: 153-162, 1961.
- CLOUGH, B.; ATILOLA, F.A.; HEALY, N. *et al.* - *Plasmodium falciparum* lacks sialidase and trans-sialidase activity. **Parasitology**, 112: 443-449, 1996.
- FORESTO, P.G.; D'ARRIGO, M.; FILIPPINI, F.R. *et al.* - Estudio de parámetros hemorreológicos en hipertensión esencial. **Rev. Fed. arg. Cardiol.**, 31: 69-73, 2002.
- HEMPHILL, A. & ROSS, C.A. - Flagellum-mediated adhesion of *Trypanosoma congolense* to bovine aorta endothelial cells. **Parasit. Res.**, 81: 412-420, 1995.
- LEBENSOHN, N.; D'ARRIGO, M.; FORESTO, P. & VALVERDE, J. - Blood stored: study of erythrocyte charge surface properties. **Vox sang.**, 93(suppl. 1): 104, 2007.

11. LEBENSOHN, N.; RE, A.; RADCLIFFE, S. *et al.* - Alterations of the erythrocyte superficial charge in glycosylated samples *in vitro*. **Biocell**, **31**(1): 137, 2007. (Abstract 90).
12. LEBENSOHN, N.; RE, A.; FILIPPINI, F. *et al.* - Impacto hemorreológico de la carga aniónica eritrocitaria en vasculopatías. **Medicina (Buenos Aires)**, **67**(suppl. III): 94, 2007.
13. PONCE DE LEÓN, P.; VALVERDE, J. & ZDERO, M. - Preliminary studies on antigenic mimicry of *Ascaris lumbricoides*. **Rev. Inst. Med. trop. S. Paulo**, **42**: 295-296, 2000.
14. RE, A.; LEBENSOHN, N.; VALVERDE, J.; D'ARRIGO, M. & FORESTO, P. - **Modelización *in vitro* de las alteraciones de carga aniónica eritrocitaria en diabéticos**. In: IX CONGRESO. XXVII REUNIÓN ANUAL DE LA SOCIEDAD DE BIOLOGIA DE ROSARIO, Rosario, Noviembre 2007. Rosario, UNR Editora, 2007. **Publicación Anual 2007**, p. 14.
15. REH, G.; NERLI, B. & PICÓ, G. - Isolation of alpha-1-antitrypsin from human plasma by partitioning in aqueous biphasic systems of polyethyleneglycol-phosphate. **J. Chromatogr.**, **780**: 389-396, 2002.
16. SHORE GARCÍA, L. & ASH, L. - **Diagnóstico parasitológico**. Buenos Aires, Editorial Médica Panamericana, 1983.
17. WACKERLY, D.; MENDENHALL, W. & SCHEAFFER, R. - **Estadística matemática con aplicaciones**. México, Thomson Learning, 2003.

Received: 8 November 2008

Accepted: 7 April 2009